

IS HEART DISEASE STILL THE NUMBER ONE KILLER IN THE POPULATION?

Kimberly Lloyd M.S.

Heart disease in the United States represents a continuing crisis of epidemic proportions, with nearly 960,000 individuals dying from heart disease and stroke each year.¹ The statistics show one out of four will have heart disease.

- Cardiovascular disease (CVD) has been the leading cause of death in this country every year since 1900, with the exception of 1918, the year of the great influenza epidemic.¹
- Heart failure is an epidemic in healthcare worldwide, now including Asia.²
- Although often perceived as a disease of older age, approximately 50 percent of CVD diagnoses and 15 percent of CVD deaths are in patients under 65 years of age.¹

Of greater significance, many young adults with no clinical evidence of CVD have two or more risk factors that predispose them to subsequent clinical events and death over the course of several decades. In apparently healthy young people, these risk factors usually go unrecognized and untreated.² Young people are not as inclined to eat much fish, regular or high omega-fatty fish.

Some types of fish contain high levels of mercury, PCBs (polychlorinated biphenyls), dioxins and other environmental contaminants. For instance, canned tuna is not recommended more than twice weekly, because of its mercury content.

The American Heart Association recommends eating fatty fish at least two times a week to obtain partial health benefits from omega-3s and omega-6s and recommends taking an omega supplement to reduce risk of heart disease. Many studies now indicate benefits of consuming one to four fish oil capsules per day.

The chart on the right shows the amount of EPA + DHA (omega-3 fatty acids) contained in a serving of various fish and shellfish. The daily use of a dietary supplement is the easiest and most logical solution. OMEGA+++ comes to the rescue with omega-3 fatty acids sourced exclusively from the *Engraulis Ringens*, a member of the anchovy family that has the highest occurring ratios of EPA and DHA of any fish species, from the pristine waters off the coast of Peru. LifePharm Global takes it a step further by purifying the fish oil to remove any possible contaminants.

According to the most recent statistics, if all forms of major CVD were eliminated, life expectancy would rise almost seven years.¹ Many risk factors are modifiable, and compelling evidence indicates that a major proportion of the current CVD burden is either totally or partially preventable. Data from the Nurse's Health Study, for example, suggest that women who maintain a desirable body weight, consume a healthy

Type of fish	Content of Omega oils (EPA + DHA) in 3 oz. fish (grams)
Tuna, Light, canned in water, drained	0.26
Sockeyed Salmon	0.68
Flounder/Sole	0.42
Halibut	0.4 - 1.0
Atlantic Cod	0.24
Shrimp	0.27
Crab	0.35
OMEGA+++ 2 soft-gel capsules	0.70

IS HEART DISEASE STILL THE NUMBER ONE KILLER IN THE POPULATION?

Kimberly Lloyd M.S.

diet, exercise regularly, avoid smoking, and drink alcohol in moderation reduce their risk of CVD by 84 percent. Yet only 3 percent of the women studied followed these recommendations, which points to the need for more effective public health messages, and changes in the healthcare system that promote prevention.

Can Omega 3 Fatty Acid Supplementation Truly Support Heart Health?

In a recent survey, three quarters of the cardiologists who recommended supplements for heart health said that if they had to choose one dietary supplement to specifically support cardiovascular health, they would select omega-3 fatty acids. New research also supports their decision. A multi-ethnic study, published in the Journal of the American Heart Association, suggests that omega-3s not only reduce cardiac threats, but do so by as much as 51 percent.

The U.S and Canadian governments have now selected omega-3s as nutrients worthy of the in-depth review process for setting a Dietary Reference Intake (DRI). Up until now, neither the FDA nor the Institute of Medicine has gone through the elaborate process of establishing any nutrient content claims for DHA and EPA. It is good to know that our health organizations are going forward for particular nutritional ingredients now found to be effective towards heart health.

The largest study that first ascertained there could be a strong association for omega fatty acid supplementation was designed to evaluate 11,324 patients who were receiving conventional heart medication after experiencing a recent heart attack. The purpose of the study was to observe if fish oil supplementation could support the prevention of a second heart attack.

- After 3.5 years of follow-up, the group that received 850 mg fish oil containing EPA and DHA experienced a 15 percent reduction in death, nonfatal heart attack, and nonfatal stroke.
- There was a 20 percent reduction in all-cause mortality and a 45 percent reduction in sudden death compared with the control group that did not receive any supplements.
- Triglycerides decreased by 4 percent and HDL cholesterol (good cholesterol) levels increased by 2.5 percent after six months in the omega-3 fatty acid treatment groups compared with controls.

Two commonly overlooked secret weapons for heart health

The exceptional OMEGA+++ formula contains other ingredients that are also beneficial for heart health.

- CoQ10 is a hardworking ingredient. CoQ10 supplementation is good for the heart to keep the energy reactions optimal. Here's why: The heart is a highly active muscle, which works continuously. Highly functioning cells contain more mitochondria than regular cells. They work much harder and need more cellular energy; the mitochondria provide this additional energy. CoQ-10 is a cofactor found in the mitochondria, which keeps the energy reactions functioning at their highest capacity.⁷
- Another ingredient in the OMEGA+++ formula that is shown to reduce the risk of heart disease is vitamin K2. Research is showing an alarming deficiency in the population for vitamin K2. Typical Western diets only provide about 10 percent of what we need to reach a recommended daily allowance for vitamin K2.^{5,6} The vitamin K2 in OMEGA+++ is from a uniquely fermented soybean product called Natto. It is highly bioavailable and this assures that it will be absorbed so the body can utilize it.

IS HEART DISEASE STILL THE NUMBER ONE KILLER IN THE POPULATION?

Kimberly Lloyd M.S.

A study selected 244 healthy postmenopausal women to receive either placebo or vitamin K2 for three years. After three years of vitamin K2 supplementation, there was a 50 percent improvement in arterial health parameters compared to placebo. The researchers concluded that long-term use of vitamin K2 supplements improved arterial stiffness in healthy postmenopausal women, especially in women having a high arterial stiffness.⁶

Do you want to live long or risk premature death?

Common sense dictates that people of all ages should eat plenty of fresh fruit and vegetables, decrease consumption of hydrogenated fats and sugar, exercise regularly and get plenty of rest in order to maintain heart health and overall health. There is more than enough evidence that good behavior alone is not enough for many segments of the population, as heart disease continues to kill at a startling rate. OMEGA+++ provides a simple and cost-effective way to supplement imperfect diets and promote heart health. The question is not "why take OMEGA+++" but "why wouldn't anyone who loves life not take OMEGA+++"?



References:

1. Robert O. Bonow, MD. Editorial Primary Prevention of Cardiovascular Disease A Call to Action *Circulation*. 2002; 106: 3140-3141 doi: 0.1161/01.CIR.0000048067.86569.E1
2. Shimokawa H, Miura M, Nochioka K, Sakata Y. Heart failure as a general pandemic in Asia. *Eur J Heart Fail*. 2015 Jul 29. doi: 10.1002/ejhf.319. [Epub ahead of print]
3. Schocken DD1, Benjamin EJ, Fonarow GC, Krumholz HM, Levy D, Mensah et al. *Circulation*. 2008 Prevention of heart failure: a scientific statement from the American Heart Association Councils on Epidemiology and Prevention, Clinical Cardiology, Cardiovascular Nursing, and High Blood Pressure Research; Quality of Care and Outcomes Research Interdisciplinary Working Group; and Functional Genomics and Translational Biology Interdisciplinary Working Group. May 13;117(19):2544-65. doi: 10.1161/CIRCULATIONAHA.107.188965. Epub 2008 Apr 7.
4. Penny M. Kris-Etherton, PhD, RD; William S. Harris, PhD; Lawrence J. Appel, MD, MPH; for the Nutrition Committee AHA Scientific Statement Fish Consumption, Fish Oil, Omega-3 Fatty Acids, and Cardiovascular Disease. *Circulation*. 2002; 106: 2747-2757 doi: 10.1161/01.CIR.0000038493.65177.94.
5. Geleijnse JM et al. "Dietary intake of Menaquinone is associated with reduced risk of coronary heart disease: The Rotterdam Study." *Journal of Nutrition* 2002 Nov. 134;(11) 3100-3105.
6. Knapen MH, et al. "Menquinone-7 supplementation improves arterial stiffness in healthy postmenopausal women. A double-blind randomised clinical trial" *Thrombosis and haemostasis* 2015 May; 113(5): 1135-1144. DOI 10.11601 TH 14-08
7. Mortensen SA1, Vadhanavikit S, Muratsu K, Folkers K Coenzyme Q10: clinical benefits with biochemical correlates suggesting a scientific breakthrough in the management of chronic heart failure. *Int J Tissue React*. 1990;12(3):155-62.